

MINISTRY OF EDUCATION, SINGAPORE
in collaboration with
CAMBRIDGE ASSESSMENT INTERNATIONAL EDUCATION
General Certificate of Education Advanced Level
Higher 2



CANDIDATE
NAME

CENTRE
NUMBER

S				
---	--	--	--	--

INDEX
NUMBER

--	--	--	--

MATHEMATICS

9758/01

Paper 1

October/November 2020

3 hours

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF26)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name on the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE ON ANY BARCODES.

Answer **all** the questions.

Write your answers in the spaces provided in the Question Paper.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

You are expected to use an approved graphing calculator.

Unsupported answers from a graphing calculator are allowed unless a question specifically states otherwise.

Where unsupported answers from a graphing calculator are not allowed in a question, you are required to present the mathematical steps using mathematical notations and not calculator commands.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **22** printed pages and **2** blank pages.



Singapore Examinations and Assessment Board



Cambridge Assessment
International Education



1 A plane π_1 contains two vectors $\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ -5 \\ -2 \end{pmatrix}$.

(i) Find a vector normal to π_1 .

[2]

A plane π_2 has equation $4x + 5y - 6z = 0$.

(ii) Find the acute angle between π_1 and π_2 .

[2]



DO NOT WRITE IN THIS MARGIN



- 2 A curve has equation $\frac{x^2}{1+x^2} + \frac{y^2}{1+y^2} = x^3y^5$. Find the equation of the tangent to the curve at the point (1, 1). Give your answer in the form $ax + by = c$, where a , b and c are integers. [6]





3 It is given that $f(x) = \ln(1 + \sin 3x)$.

(i) Show that $f''(x) = \frac{k}{1 + \sin 3x}$, where k is a constant to be found.

[3]

DO NOT WRITE IN THIS MARGIN





(ii) Hence find the first three non-zero terms of the Maclaurin expansion of $f(x)$.

[4]

$$f(x) = \frac{1}{1-x^2} = (1-x^2)^{-1}$$

$$= 1 + x^2 + x^4 + \dots$$





4 Do not use a calculator in answering this question.

Three complex numbers are $z_1 = 1 + \sqrt{3}i$, $z_2 = 1 - i$ and $z_3 = 2(\cos \frac{1}{6}\pi + i \sin \frac{1}{6}\pi)$.

(i) Find $\frac{z_1}{z_2 z_3}$ in the form $r(\cos \theta + i \sin \theta)$, where $r > 0$ and $-\pi < \theta \leq \pi$.

[4]





A fourth complex number, z_4 , is such that $\frac{z_1 z_4}{z_2 z_3}$ is purely imaginary and $\left| \frac{z_1 z_4}{z_2 z_3} \right| = 1$.

(ii) Find the possible values of z_4 in the form $r(\cos \theta + i \sin \theta)$, where $r > 0$ and $-\pi < \theta \leq \pi$. [3]

... (b) ...

... (c) ...





5 (a) Given that \mathbf{a} and \mathbf{b} are non-zero vectors such that $\mathbf{a} \times \mathbf{b} = \mathbf{b} \times \mathbf{a}$, find the relationship between \mathbf{a} and \mathbf{b} . [2]

(b) The points P , Q and R have position vectors \mathbf{p} , \mathbf{q} and \mathbf{r} respectively. The points P and Q are fixed and R varies.

(i) Given that \mathbf{q} is non-zero and $(\mathbf{r} - \mathbf{p}) \times \mathbf{q} = \mathbf{0}$, describe geometrically the set of all possible positions of the point R . [3]





- (ii) Given instead that $\mathbf{r} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$, $\mathbf{p} = \begin{pmatrix} -1 \\ 2 \\ 4 \end{pmatrix}$, $\mathbf{q} = \begin{pmatrix} 3 \\ -5 \\ 2 \end{pmatrix}$ and that $(\mathbf{r} - \mathbf{p}) \cdot \mathbf{q} = 0$, find the relationship between x , y and z . Describe the set of all possible positions of the point R in this case. [4]





6 The complex number z satisfies the equation

$$z^2(2 + i) - 8iz + t = 0,$$

where t is a real number. It is given that one root is of the form $k + ki$, where k is real and non-zero.

Find t and k , and the other root of the equation.

[8]





6 [Continued]





7 Do not use a calculator in answering this question.

It is given that $f(x) = 2 - \sin 4x$.

(i) Find $\int f(x) dx$. [1]

(ii) Find the exact value, in terms of π , of $\int_0^{\frac{1}{2}\pi} x f(x) dx$. [4]





(iii) Find the exact value, in terms of π , of $\int_0^{\frac{1}{2}\pi} (f(x))^2 dx$.

[4]





8 (a) The 1st term of an arithmetic series is 4 and the 5th term is 10.

(i) Find the 30th term of this series.

[2]

(ii) Find the sum of the 21st term to the 50th term inclusive of this series.

[3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN





(b) The 1st term of a geometric series is 4 and the 5th term is 1.6384 where the common ratio is positive.

(i) Find the sum to infinity of this series. [2]

(ii) Given that the sum of the first n terms is greater than 19.6, show that

$$0.8^n < 0.02.$$

Hence find the smallest possible value of n . [5]





- 9 (i) By considering the gradients of two lines, explain why $\tan^{-1}(2) - \tan^{-1}\left(-\frac{1}{2}\right) = \frac{1}{2}\pi$. [1]

The curves C_1 and C_2 have equations $y = \frac{1}{x^2 + 1}$ and $y = \frac{k}{3x + 4}$ respectively, where k is a constant and $k > 0$.

- (ii) Find the set of values of k such that C_1 and C_2 intersect. [3]





It is now given that $k = 2$.

(iii) Sketch C_1 and C_2 on the same graph, giving the coordinates of any points where C_1 or C_2 cross the axes and the equations of any asymptotes. [3]

(iv) Find the exact area of the region bounded by C_1 and C_2 , simplifying your answer. [5]





10 Scientists are investigating the effect of disease on the number of sheep on a small island. They discover that every year the death rate of the sheep is greater than the birth rate of the sheep. The difference every year between the death rate and the birth rate for the population of sheep on the island is 3%. The number of sheep on the island is P at a time t years after the scientists begin observations.

(i) Write down a differential equation relating P and t .

[2]

(ii) Solve this differential equation to find an expression for P in terms of t . Explain what happens to the number of sheep if this situation continues over many years.

[4]





The scientists import sheep at a constant uniform rate of n sheep per year. (The difference every year between the death rate and the birth rate remains at 3%.)

(iii) Write down a differential equation to model the new situation. [2]


(iv) Solve the differential equation to find an expression for P in terms of t and n . [4]

(v) Given that the number of sheep settles down to 500 after many years, find n . [2]





- 11 In sport science, studies are made to optimise performance in all aspects of sport from fitness to technique.



In a game, a player scores 3 points by carrying the ball over the scoring line, shown in the diagram as XY . When a player has scored these 3 points, an extra point is scored if the ball is kicked between two fixed vertical posts at C and D . The kick can be taken from any point on the line AB , where A is the point at which the player crossed the scoring line and AB is perpendicular to XY .

The distance CD is 4 m; XC is equal to DY ; the point A is a distance a m from C and A lies between X and C . The kick is taken from the point K , where AK is x m. The angle CKD is θ (see diagram).





(i) By expressing θ as the difference of two angles, or otherwise, show that

$$\tan \theta = \frac{4x}{x^2 + 4a + a^2} \quad [3]$$

(ii) Find, in terms of a , the value of x which maximises $\tan \theta$, simplifying your answer. Find also the corresponding value of $\tan \theta$. (You need not show that your answer gives a maximum.) [3]





11 [Continued]

The point corresponding to the value of x found in part (ii) is called the optimal point. The corresponding value of θ is called the optimal angle.

(iii) Explain why a player may decide not to take the kick from the optimal point. [1]

(iv) Show that, when θ is the optimal angle, $\tan KDA = \sqrt{\frac{a}{4+a}}$. Find the approximate value of angle KDA when a is much greater than 4. [3]

(v) It is given that the length of the scoring line XY is 50 m. Find the range in which the optimal angle lies as the location of A varies between X and C . [2]

